

NOTICE CONCERNING COPYRIGHT RESTRICTIONS

This document may contain copyrighted materials. These materials have been made available for use in research, teaching, and private study, but may not be used for any commercial purpose. Users may not otherwise copy, reproduce, retransmit, distribute, publish, commercially exploit or otherwise transfer any material.

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material.

Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specific conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

This institution reserves the right to refuse to accept a copying order if, in its judgment, fulfillment of the order would involve violation of copyright law.

The Reykjanes Geothermal System: Alteration Mineralogy and Fluid-Rock Interaction in an Icelandic High Temperature System

Naomi Marks, Peter Schiffman, Robert Zierenberg

Department of Geology, University of California-Davis, Davis, CA, USA

marks@geology.ucdavis.edu

Keywords

Iceland, basalt, geothermal systems, hydrothermal alteration, Sr isotopes

ABSTRACT

The Reykjanes geothermal system is a seawater-recharged hydrothermal system that appears to be analogous to seafloor hydrothermal systems, and for this reason is a useful proxy for seafloor vents. During the Pleistocene, the system was dominated by meteoric water recharge, and fluid composition at Reykjanes has evolved through time as a result of changing proportions of meteoric water influx as well as differing pressure and temperature conditions. The purpose of this study is to characterize secondary mineralization, degree of metasomatic alteration, and bulk composition of cuttings from well RN-17 from the Reykjanes geothermal system. The progressive hydrothermal alteration sequence observed with increasing depth results from reaction of geothermal fluids with the basaltic host rock. Alteration phase textures of drill cuttings reveal alteration style to be strongly dependant on protolith. Hyaloclastites are intensely altered with calc-silicate alteration assemblages comprising calcic hydrothermal plagioclase, grandite garnet, prehnite, epidote, hydrothermal clinopyroxene, and titanite. In contrast, crystalline basalts and intrusives display a range in alteration intensity from essentially unaltered to pervasive and nearly complete albitization of igneous feldspar and uralitization of clinopyroxene. $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic values of alteration phases including epidote range from 0.7037 to 0.7076 and indicate isotopic exchange with seawater Sr.

Introduction

The Reykjanes geothermal field, located on the landward extension of the Mid-Atlantic Ridge, is the location of a 3 km deep drill-hole developed by the Iceland Deep Drilling Project to investigate supercritical fluids as a potential energy resource and to explore geologic processes in actively spreading margins (Figure 1). In order to evaluate the nature and extent of rock-

fluid interactions and the applicability of Reykjanes as a proxy to oceanic spreading margins, it is necessary to understand the source and composition of hydrothermal fluids within the system. The composition and salinity of geothermal fluids at Reykjanes resemble evolved seawater, suggesting that subsurface conditions at Reykjanes may be analogous to sea-floor black smokers. The high temperature reaction zone that is presumed to control the composition of the hydrothermal fluid is interpreted to occur deeper than the present depth of drilling, which reached just over 3 km in the well RN-17, during the initial stage of the Iceland Deep Drilling Project. Fluid composition at Reykjanes has evolved through time as a result of changing proportions of meteoric water influx as well as differing pressure and temperature conditions imposed by glaciation (Arnorsson, 1995; Fridleifsson et al.,




Figure 1. Map of the Reykjanes Peninsula showing major faults, fractures, flows, surface alteration, and the location of IDDP well of opportunity RN-17 (map adapted from Saemundsson 2000, Karlsdottir 1998 and Fridleifsson et al. 2000, and Franzson et al., 2002).

2003). Reaction of geothermal fluids with the basaltic host rock has resulted in a progressive hydrothermal alteration sequence with increasing depth (Fridleifsson et al, 2005; Franzson et al, 2002; Lonker et al., 1993). The assemblage of greenschist facies alteration minerals implies that temperatures reached at least 250°C at depths as shallow as 350 m. This requires hydrostatic pressures that exceed the present-day boiling point to depth curve, and therefore must record alteration at higher fluid pressures during the Pleistocene.

Methods

Cuttings were collected from RN-17 at 2m intervals from the entire depth of the well for lithologic logging. Cuttings were inspected with a binocular microscope as well as in 34 thin sections. Petrographic and X-ray diffraction studies were carried out on these cuttings at ISOR, the Icelandic Geosurvey.

Backscattered electron imaging (BSE) and mineral compositional analyses were performed at UC Davis on cuttings selected at 50 m intervals. Cuttings were mounted on doubly polished, carbon coated thin sections and analyzed using a Cameca SX-

100 electron microprobe. Quantitative wavelength-dispersive analyses of amphibole, epidote, chlorite, garnet, and prehnite were performed at 15 keV and 10 nA beam current using an approximately 1 μm diameter beam. Feldspar analyses were performed at 15 keV and 10 nA beam current using an approximately 5 μm beam.

Sr isotopic measurements were made on secondary phases by laser ablation multicollector inductively coupled plasma mass spectrometry at UC Davis.

Results

Well lithology is presented in Figure 2. Samples from the deepest part of Reykjanes well RN-17 include greenschist and amphibolite facies assemblages, suggesting seawater penetration into a part of the system that is close to the high temperature reaction zone. Electron microprobe studies of drill cuttings reveal intense alteration of hyaloclastites with calc-silicate alteration assemblages comprising calcic hydrothermal plagioclase, grandite garnet, prehnite, epidote, hydrothermal clinopyroxene, and titanite. In contrast, crystalline basalts and intrusives display a wide range in alteration intensity from essentially unaltered to pervasive and nearly complete replacement of feldspar and pyroxene.

Anhydrite is widespread in shallow portions of the Reykjanes system to about 1500 m, and after a depositional hiatus appears again around 2650 m depth. This may suggest either a transient incursion of cold seawater into the system at depth, or possibly a more permanent cooling at depth in the system. A zone of amphibolite facies contact metamorphism has also been observed at this depth (Figure 2), indicating temperatures as high as 600°C (Marks et al., 2008), and the reappearance of anhydrite could possibly indicate heating and oxidation of previously precipitated sulfides.

⁸⁷Sr/⁸⁶Sr values of individual epidote grains were typically 0.7045-0.7050, but ranged as high as 0.7073 in individual epidote grains (Figure 3).

Discussion

Epidote occurs in virtually all geothermal systems in which present or calculated paleotemperatures exceed 200-250°C (Bird et al., 1984). Epidote is widely distributed throughout the RN-17 samples as veins and vug fillings, replaces hyaloclastite and the interstitial matrix of basalt samples, and is also an alteration product of primary plagioclase. It occurs preferentially in the more reactive hyaloclastite and is associated with prehnite, chlorite, quartz, zeolites, calcite, amphibole, and garnet. Strontium isotopic ratios in alteration phases such as epidote indicate a shift from basaltic values towards the seawater Sr values suggesting penetration of seawater Sr to great depth within the system.

A New Alteration Facies

Alteration phases indicative of a transition from Epidote-Actinolite alteration to alteration at amphibolite conditions are not readily observed during routine cuttings logging, but were documented during BSE petrographic investigation and the mineral chemistry of these phases is reported in following sections.




Figure 2. Lithostratigraphy, alteration zones, and distribution of secondary minerals (below the smectite zone) with depth in RN-17 (modified after Fridleifsson et al., 2005).




Figure 3. $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic ratios measured in epidote from RN-17. Bars indicate 2σ error.

We define a new alteration facies, the amphibole zone, based on the occurrence of the high temperature minerals hornblende-clinopyroxene-plagioclase. The facies boundary is transitional with the first occurrence of these minerals at 2350 meters, which are locally developed within the Epidote-Actinolite Facies to the bottom of the hole.

Effect of Protolith on Intensity of Alteration

The nature of the protolith strongly influences the intensity of alteration in this rock dominated hydrothermal system. The hyaloclastites, tuffs, and sediments are more susceptible to alteration than the crystalline basalts as a result of their greater reactivity, greater percentage of glass, and abundant vesicles. Throughout RN-17, hyaloclastites are highly altered and host most of the hydrous calc-silicates, plus calcite and chlorite. Anhydrite and sulfide minerals are almost exclusively hosted in hyaloclastite.

In contrast, the degree of alteration in basalts is highly variable. While basalts range from essentially unaltered to thoroughly albitized and uraltized, unaltered basalts are frequently found in the same cutting interval as pervasively altered hyaloclastites. The most altered basalts host abundant amphibole up to magnesiohornblende in composition, as well as secondary orthoclase feldspar and, rarely, hydrothermal anorthite. The altered basalts are almost exclusively from cuttings intervals dominated by pil-

low basalts, which seem to alter more readily due to cracking, vesiculation, and a higher proportion of glass that enhances fluid ingress. Crystalline basalts are minimally altered in general. Both hyaloclastite and crystalline basalt cuttings host secondary veins, which typically comprise quartz, calcite, epidote and, rarely, hydrothermal plagioclase. The formation of calc-silicate minerals and aluminous amphiboles requires release and mobilization of aluminum during alteration (Schiffman, 1995). Formation of these alteration phases in crystalline basalts therefore requires extensive alteration of plagioclase to albite. In contrast, in hyaloclastite or pillow basalts, alteration of glass can provide a local source of aluminum and may contribute to the more extensive alteration observed in these lithologies.

Sr Isotopic Ratios

The Sr isotopic ratios of alteration minerals are shifted from basaltic values (0.7030-0.7034; O'Nions and Grönvold, 1973; Sun and Jahn, 1975) toward seawater values (0.70916; Palmer and Edmond, 1989). Epidote samples from the Reykjanes system have significantly higher Sr isotopic ratios than analogous freshwater dominated geothermal systems at Nesjavellir and Krafla. This suggests that seawater Sr is able to penetrate deep within the geothermal system, and that seawater Sr has been able to exchange with secondary phases throughout the system.

Conclusions

Samples from the Iceland Deep Drilling Program (IDDP) well of opportunity RN-17 have provided a basis for the study of the geochemistry of alteration within the Reykjanes system. We have found the intensity of alteration and secondary mineralization to be directly related to the texture of the protolith. Specifically, high porosity, glassy fragments are more susceptible to alteration, and holocrystalline basalts typically display minimal alteration. We have used electron microprobe and BSE methods to define the amphibole zone, a new alteration facies within RN-17. This zone indicates higher temperatures and pressures than had previously been suggested for the base of this well. Finally, $^{87}\text{Sr}/^{86}\text{Sr}$ ratios within alteration minerals are significantly shifted toward seawater values within the system, and therefore confirm the deep penetration of seawater into the Reykjanes system.

References

- Arnorsson, S. (1995). Geothermal systems in Iceland: structure and conceptual models-I. High-temperature areas. *Geothermics*, 24(5):561-602.
- Bird, D., Schiffman, P., Elders, W., Williams, A., and McDowell, S. (1984). Calc-silicate mineralization in active geothermal systems. *Economic Geology*, 79:671-695.
- Franzson, H., Thordarson, S., Bjornsson, G., Gudlaugsson, S., Richter, B., Fridleifsson, G., and Thorhallsson, S. (2002). Reykjanes high-temperature field, SW-Iceland. Geology and hydrothermal alteration of well RN-10. Workshop on Geothermal Reservoir Engineering, 27:233-240.
- Fridleifsson, G., Arnannsson, H., Arnason, K., Bjarnason, I., and Gislason, G. (2003). Part I: Geosciences and Site Selection. In Fridleifsson, G., editor, Iceland Deep Drilling Project, Feasibility Report, volume OS-2003-007, page 104. Orkustofnun Report.

- Fridleifsson, G. and Elders, W. (2005). The Iceland Deep Drilling Project: a search for deep unconventional geothermal resources. *Geothermics*, 34(3):269–285.
- Lonker, S., Franzson, H., and Kristmannsdottir, H. (1993). Mineral-fluid interactions in the Reykjanes and Svartsengi geothermal systems, Iceland. *American Journal of Science*, 293(7):605–670.
- Marks, N., Schiffman, P., Zierenberg, R., and Franzson, H. (2008). The Iceland Deep Drilling Project: (III) Evidence for amphibolite grade contact metamorphism in an active geothermal system. In American Geophysical Union, Fall Meeting 2008, abstract# V41B-2071.
- O’Nions, R. and Gronvold, K. (1973). Petrogenetic relationships of acid and basic rocks in Iceland: Sr-isotopes and rare-earth elements in late and post-glacial volcanics. *Earth and Planetary Science Letters*, 19(4):397–409.
- Palmer, M. and Edmond, J. (1989). The strontium isotope budget of the modern ocean. *Earth and Planetary Science Letters*, 92(1):11–26.
- Schiffman, P. and Day, H. (1995). Low-grade metamorphism of mafic rocks. Geological Society of America.
- Sun, S.-S. and Jahn, B. (1975). Lead and strontium isotopes in post-glacial basalts from Iceland. *Nature*, 255(5509):527–530.